



## Macrophyte Occurrence in Response to Anthropogenic Pressure in Reservoir

Carla Denise Tedesco <sup>1</sup>

Claudia Petry <sup>2</sup>

Edson Campanhola Bortoluzzi <sup>3</sup>

Alfredo Castamann <sup>4</sup>

### ABSTRACT

The objective of this study was to evaluate the effect of anthropogenic land use on the community of aquatic plants in the vicinity of a consolidated artificial reservoir. Two landscape units (LU) were outlined adjacent to a more than 50-year-old artificial reservoir, of which one (LU1) had a low and the second (LU2) a high human impact. In total, we found 20 plant species. Only two species were common to UP1 and UP2. The species diversity in LU1 (n=18) was greater than in LU2 (n=4), and plant distribution more even. Phosphorus, potassium, organic matter, calcium, and magnesium contents were lower in LU2. It can be concluded that the embankment induced the occurrence of more macrophyte species and plants, due to the soil conservation in the reservoir surroundings. The low human impact in this area allowed a greater diversity of native macrophytes. The intensification of land use around the reservoir impoverishes the soil, preventing the development of a diversified community of macrophyte.

**Keywords:** Aquatic Plants; Wetland; Conservation.

---

<sup>1</sup> Doutorado em Agronomia pela Universidade de Passo Fundo, UPF, Brasil. Docente na Universidade de Passo Fundo, UPF, Brasil. [tedesco@upf.br](mailto:tedesco@upf.br)

<sup>2</sup> Doutorado em Geografia pela Université Paris 1 Pantheon-Sorbonne, PARIS 1, França. Docente na Universidade de Passo Fundo, UPF, Brasil. [petry@upf.br](mailto:petry@upf.br)

<sup>3</sup> Doutorado em Science Du Sol Et Environnement pelo Institut National Agronomique Paris Grignon, INA P-G, França. Docente na Universidade de Passo Fundo, UPF, Brasil. [edsonb@upf.br](mailto:edsonb@upf.br)

<sup>4</sup> Doutorado em Agronomia pela Universidade de Passo Fundo, UPF, Brasil. Docente na Universidade Federal da Fronteira Sul, UFFS, Brasil. [alfredo.castamann@uffs.edu.br](mailto:alfredo.castamann@uffs.edu.br)

In Brazil, 87.1% of the power generation capacity is hydroelectric (OECD, 2015). The embankment of rivers leads to the flooding of vast areas on plains with small slopes or even on plateaus, and is of fundamental to sustain the production of this type of electricity (Kelman et al. 2006).

Due to the flooding, damming leads to a series of environmental changes in the surrounding areas, resulting in the replacement of the original plant community by species adapted to a changing water level (amphibious), or those of aquatic environments (floating and submerged species) (Mukhopadhyay & Dewanji 2005, Peintinger et al. 2007). The chemical and physical water quality is changed with the increase of nitrogen and phosphate compounds and with turbidity (Wetzel 2001, Aitkenhead-Peterson et al. 2003). Also, the sediments introduced from adjacent areas represents another factor of reduction of the wetland quality and lifetime of the proper reservoir for energy generation. Aside from hampering the organoleptic quality of drinking water, these processes affect power generation. The water quality of the reservoir is also a consequence of the water quality in the neighboring areas, which were modified by the construction of the reservoir, with deforestation and the removal of natural soil horizons, affecting the soil functionality (Zhao et al. 2014; Navarro-Ortega et al. 2015). This functionality of the soil is understood as the barrier effect of the areas surrounding the reservoir, which can reduce or prevent the arrival of nutrients, pollutants and sediments from terrestrial systems. Thus, the environmental quality in the flooded area, which defines the quality of soil and biota in these ecosystems, depends on the level of anthropogenic pressure in the surroundings of the reservoir (Banach et al. 2009, Sass et al. 2010, Maëmets et al. 2010).


Brazil is home to about 980 macrophyte species (Chambers et al. 2008) in naturally flooded areas. The Pantanal of Mato Grosso comprises 280 species (Pott et al. 2011) and the upper Rio Paraná floodplain over 150 species (Ferreira et al. 2011). In southern Brazil, lagoons and coastal wetlands have varying species richness. In studies in this region, Pereira et al. (2012) recorded 43 species; Maltchik et al. (2011) 59 species; Spellmeier et al. (2009) 145 species; and Rolon et al. (2011) 176 species. Due to the richness of species and life forms of macrophytes, they play a key role in post-flooded environments and can serve as an indicator parameter of their quality.

We tested the hypothesis that flooding provides conditions for the development of new macrophyte communities, but that the species occurrence and frequency will depend on the subsequent land use in the areas surrounding the artificial reservoir. In this sense, the purpose of the study was the comparison of macrophyte communities in two landscape units, adjacent to an artificial reservoir, with different anthropogenic pressure. To this end, the occurrence of macrophyte species was investigated to relate their characteristics and distribution in the landscape units, as well as soil and water aspects.

## MATERIAL AND METHODS

The reservoir of Capiguí, located in southern Brazil (Figure 1), was flooded in 1956 (CEEE, 2011). The soils are classified as Nitisols and Ferralsols (at higher altitudes) and poorly drained Gleysols (IUSS 2014).

**Figure 1.** A: Latin America, southern Brazil. B: State of Rio Grande do Sul and Passo Fundo, municipality where the reservoir is located; C: Reservoir of Capiguí; D: Transect perpendicular to the margin of Landscape Unit 2; E: Overview of Landscape Unit 1



Fonte: Os Autores.

The climate is humid subtropical (Cfa according to Köppen), with no dry season, temperature of the warmest month above, mean annual temperature 17.5 °C, and mean annual rainfall 1787.8 mm. The relative air humidity is on average 72% (Cunha 1997).

The reservoir covers an area of 6.27 Km<sup>2</sup> and has a perimeter of 43.43 km. Prior to flooding, the region was covered with native grasslands, forest fragments and family farming. The current land use in the 100 m around the reservoir consists of field/pasture or agriculture (52%); forest cover with varying degrees of conservation (40%); residential areas (6%) and forestry (2%) (CEEE 2011).

### **CHOICE OF ADJACENT AREAS (LANDSCAPE UNITS)**

Fifty-two years after the reservoir had been constructed, two landscape units were selected on farms with the presence of aquatic plants during visits to the reservoir (Figure 1). The units were chosen at random for the presence of macrophytes, visual characterization of the wetlands and the land use history, based on ethnographic information from local residents. The landscape units are temporarily flooded, mainly due to the reservoir management for energy demand. So, the two landscape units were described as follows:

- Landscape Unit 1 (LU1- 28° 20' 30.00" S and 52° 10' 18.5" W, 634 m asl, 3.6% slope), is located on a farm and represents the control: low anthropogenic pressure, with no agricultural cultivation of annual or perennial crops nor cattle grazing within a distance of 1,000 m from the reservoir. Residences are built at a distance of at least 200 m.
- Landscape unit 2 (LU2 - 28° 19' 46.2" S and 52° 13' 56.0" W, 634 m asl and slope of 3.52%): high anthropogenic pressure, characterized by land use with frequent presence of grazing cattle. Agricultural inputs (fertilizer or lime) are not used directly nor agricultural machinery for any kind of management. Landscape unit LU2 is part of a small farm, where, in addition to dairy farming, annual corn and wheat crops are grown, and 15,000 birds are fattened in a poultry-house at a distance of more than 200 m from the flooded area.

### **SAMPLING**

In the two units (LU1 and LU2) a transect (width 1 m, length 19 m - in both unit) was delimited, perpendicular to the reservoir shores (Figure 1, letters D and E), adjacent to the flooded area, with the presence of macrophytes. Wetlands presented very homogeneous as your face, so it was considered only a transect.

The following variables were recorded in field charts for each transect, divided into one m<sup>2</sup> units, totaling 19 sample units (quadrats): presence of vascular species (macroalgae were excluded), biological form (Irgang & Gastal 1996), plant cover degree (Braun-Blanquet et al. 1979), type of plant distribution, plant appearance, and sun exposure level. Thus, the relationship of number of species per quadrat between the more and less humid environments in LU1 and LU2 can be established. The macrophytes were collected from 04 to 10 January 2009. Voucher specimens of each macrophyte species were collected for herborization, and deposited in the Herbarium Rio Grande do Sul, Passo Fundo (RSPF), of the Institute of Biological Sciences, University of Passo Fundo (ICB/UPF).

Soil and water samples were collected from each LU. Composite soil samples were collected with a cutting shovel from the 0-20 cm layer, consisting of a blend of four sub-samples of about 500 g each. Water samples were collected in a clean, sterile container from the flooded part of the two landscape units. All analytical determinations of soil chemical properties were performed in the Laboratory Soil Analysis, Faculty of Agronomy and Veterinary Medicine, University of Passo Fundo, according to the methodology described by Tedesco et al. (1995). The results of soil analysis were interpreted as proposed by the Brazilian Society of Soil Science (CQFS 2016).

### **TREATMENT OF RESULTS**

After species identification, the frequency of occurrence was recorded ( $FAI = (ni/N) \times 100$ ) in each LU, where FAI = frequency of species *i*; *ni* = number of sample units where the species was recorded; *N* = total number of sample units.

The macrophytes, recorded on the Braun-Blanquet scale were converted to the ordinal scale of van der Maarel (1979) to assess the diversity index. The Shannon-Wiener diversity index was calculated using the program Divers (version 2011), to compare the landscape units. The sampling effort was evaluated using a species accumulation curve with 1000 simulations, to evaluate the sampling efficiency (Colwell & Coddington 1994), through the EstimateS 9.1.0 software (Colwell 2013). Each area one quadrat corresponded to a sample unit. Accumulation curves were obtained using non-parametric Bootstrap estimators to confirm the validation of the sampling. The soil analyses data were discussed qualitatively and related with soil and water parameters.

### **RESULTS**

The species found in the reservoir of Capigui in the units LU1 and LU2 are listed in Table 1. Fifteen families and 20 macrophyte species were recorded in both areas. The *Onagraceae* family (3 species), *Cyperaceae*, *Poaceae* and *Polygonaceae* (2 species each), were represented best. The four families accounted for 45% of all observed macrophyte species (Table 1).

In LU1 more than half of the species ( $n = 10$ ) were considered rare (frequency of occurrence less than 12%), five species were considered common (frequency between 20 and 49%) and three constants (frequency greater than 50%). The so-called amphibious species accounted for 85% of the total and this portion consisted of procumbent herbs, with irregular distribution on predominantly clay soil, exposed to full sun, with biological forms of plants rooted in the substrate and tolerate to water stress (Table 1). Also, according to the List of flora species of Brazil (2016), 18 species were classified as native to Brazil and/or South America. *Alternanthera philoxeroides* (LU2) stood out for being present in all quadrats of LU2 and *Paspalum cf notatum* only in 21,05 % (Table 1). The difference between the diversity indices (Table 1) reinforces the disparity of the LU2 community (low richness and dominance of one species) compared to LU1 (richer in and more balanced relative abundance of species).

In both LUs, the number of species was higher farther away from the flooded environment all species found were classified as amphibious (Table 1).

Tables 2 and 3 show the results of water and soil analysis. Calcium levels in the soil were fairly different between the two landscape units. In surface waters, Ca is found at concentrations up to 15 mg L<sup>-1</sup>, mostly in the minerals calcite, dolomite and gypsum (Paron et al. 2011). Magnesium occurs most often in the minerals magnesite and dolomite, at a mean concentration of 4 mg L<sup>-1</sup> in natural waters (Clesceri et al. 1999). The Al levels were clearly below the maximum value allowed for Class I water classification in Brazil, i.e., 0.1 mg L<sup>-1</sup> (Brasil 2005). The pH values were within the range considered normal (6.0 to 9.0) for all water classes. The conductivity was similar in the two LUs and indicated a low degree of water mineralization, i.e., the concentration of dissolved minerals was low (Clesceri et al. 1999).

These results indicate that the different landscape units fall into the same texture (proportion of clay), indicating a similar chemical reactivity with the environment. Similarly, soil pH values are interpreted as very low at the studied locations. The data of other properties were interpreted differently, indicating that the presence of animals in LU2 may have contributed reduce soil fertility. In LU1, higher phosphorus, potassium, organic matter, calcium and magnesium contents were observed than in LU2. The lower soil fertility may show the most intense anthropogenic effect on LU2.

## DISCUSSION

The embankment of the river Capiguí in the 1950's changed the landscape, allowing the installation and development of macrophyte communities in the originally well-drained soils. The response in species occurrence and frequency of macrophyte communities in relation to land use was

contrasting between the two landscape units. In short, the increased human pressure reduced species richness and increased the abundance of the species *Alternanthera philoxeroides*.

**Table 1.** Family and botanical species, Number of species, Shannon diversity index frequency of occurrence (%) and description of two macrophyte communities in the surroundings of the reservoir of Capigui in two landscape units: LU1 (low anthropogenic pressure) and LU2 (high anthropogenic pressure).

FAMILY/BOTANICAL SPECIES	FREQUENCY %		SPECIES DESCRIPTION						
	LU1	LU2	BF	DSU	PA	WP	ST	SE	O
<b>AMARANTHACEAE</b>									
<i>Alternanthera philoxeroides</i> (Mart.) Griseb	5.26	100.00	amp	i, r	h,p	d,a	c	sop fs	n
<b>APIACEAE</b>									
<i>Eryngium</i> sp	36.84		amp	i	h,p	ws	c	fs	ni
<b>BORAGINACEAE</b>									
<i>Echium plantagineum</i> L.	10.53		amp	i	h,p	ws	c	fs	ni
<b>CLUSIACEAE</b>									
<i>Hypericum brasiliense</i> Choisy	26.32		amp	i	h,p	ws	c	fs	n
<b>COMMELINACEAE</b>									
<i>Commelina erecta</i> L.	5.26		amp	i	h,p	ws	c	fs	n
<b>CYPERACEAE</b>									
<i>Rhynchospora aurea</i> Vahl	57.89	5.26	amp	i	h,p	ws	c	fs	n
<i>Eleocharis sellowiana</i> Kunth	21.05		amp	i	h,p	ws	c	fs	n
<b>HALORACEAE</b>									
<i>Myriophyllum aquaticum</i> (Vell.) Verdcourt	42.11		evp	i	h,p	d,a	c	sop	n
<b>JUNCACEAE</b>									
<i>Juncus effusus</i> L.	10.53		amp	i	h,p	ws	c	fs	n
<b>LYTHRACEAE</b>									
<i>Cuphea calophylla</i> Cham. et Schldt	10.53		amp	i	h,p	ws	c	fs	n
<b>MAYACACEAE</b>									
<i>Mayaca sellowiana</i> Kunth	5.26		evp	i	h,p	d,a	c	sop	n
<b>MENYANTHACEAE</b>									
<i>Nymphoides humboldtiana</i> (Kunh) Kuntze	10.53		fl	i	h,p	d,a	c	fs	n
<b>ONAGRACEAE</b>									
<i>Ludwigia multinervia</i> Hook. & Arn.		5.26	amp	i	h,p	ws	c	fs	n
<i>Ludwigia peploides</i> (Kunth) Raven	5.26		amp	i	h,p	ws	c	fs	n
<i>Ludwigia sericea</i> (Cambess.) H. Hara	57.89		amp	i	h,p	ws	c	fs	n
<b>POACEAE</b>									
<i>Paspalum cf notatum</i>		21.05	amp	i	h,p	ws	c	fs	n
<i>Panicum schwackeanum</i> Mez	31.57		amp	i	h,p	ws	c	fs	n
<b>POLYGONACEAE</b>									
<i>Polygonum acuminatum</i> Kunth	5.26		amp	i	h,p	ws	c	fs	n
<i>Polygonum meisnerianum</i> Cham.	52.63		amp	i	h,p	ws	c	fs	n
<b>XYRIDACEAE</b>									
<i>Xyris jupicai</i> Rich	5.26		amp	i	h,p	ws	c	fs	n
<b>Number of species</b>	<b>18.00</b>	<b>4.00</b>							
<b>Shannon-Wiener diversity index</b>	<b>1.03</b>	<b>0.13</b>							

BF: Biological form: fl = floating leaves, evp = emergent vegetative part, amp = amphibious (drought-tolerant, all forms rooted in the substrate). DSU: Distribution in sample unit i = irregular, r = regular. PA: Plant Appearance: h = herb, p = procumbent. WP: presence of water: ws = waterlogged soil (saturated with water); d = water depth; a = flooded area (water deeper than 3 cm). ST: Predominant Soil texture: c = clay. SE: Sun exposure: fs = exposed to full sun, sop = shaded by other plants. O: Origin, n = native to South America, ni = no information. Characteristics recorded at the moment of sampling and the origin from consultations of literature sources and herbarium.

Fonte: Os Autores.

**BOTANICAL ANALYSIS OF SPECIES OCCURRENCE**

The most common organic form (amphibious) (Table 1), may be associated with the fluctuation of the water level of the reservoir, which is designed to accumulate water for the downstream system.. The analysis of species occurrence in the transects (LU1 and 2) shows an predominance in amphibious species (Figure 1). To survive in flooded areas, plants have to undergo a series of physiological and morphoanatomical adaptations (Evans 2003, Jackson & Colmer. 2005, Bailey-Serres & Voeselek. 2008, Colmer & Voeselek 2009, Jackson et al. 2009) to survive hypoxic or anoxic environments, or anaerobic conditions of the soil (Sairam et al. 2008). This indicates that amphibious plants are highly tolerant to variations in the water level, but are affected by waterlogged soils, that possibly occur very close to the flooded area.

The variations in richness, biological forms and type of sun exposure (Table 1) were largest in LU1. The distribution of all plants in LU1 was irregular, suggesting that intraspecific competition processes regulate the species distribution. Soil fertility was higher in this area. The lowest species richness in the area with greater human disturbance was also reported by Meyer & Francheschinelli (2010) and Galvão et al. (2011). Sass et al. (2010) pointed out that different disturbance gradients in the vicinity of ponds affect species differently by increasing or decreasing their abundance.

The Shannon diversity index was lowest for LU2 (Table 1). The difference was a result of the dominance of *A. philoxeroides* and the small number of species in this LU. Importantly, these wetlands are recent, since the embankment was constructed in the 50s, so this community is assumedly still in the succession process. In the state of Rio Grande do Sul and other Brazilian regions, the Shannon diversity index is usually above 2.0 (Boldrini et al. 2008, Alves et al. 2011, Evangelista et al. 2012).

**Table 2.** Chemical properties of water collected in the landscape units LU1 and LU2.

LANDSCAPE UNIT	ANTHROPOGENIC LEVEL	PROPERTIES				
		Ca mg L <sup>-1</sup>	Mg mg L <sup>-1</sup>	Al mg L <sup>-1</sup>	pH Um	CE μS cm <sup>-1</sup>
LU1	Low	4.87	2.5	0.01	6.9	31.5
LU2	High	6.59	2.95	0.01	6.1	30.3

Fonte: Os Autores.


**Table 3.** Soil analysis results in two areas: LU1, LU2 and low anthropogenic pressure, high anthropogenic pressure.

AREA	Cy (%)	pH	SMP Ind.	P mg dm <sup>-3</sup>	K mg dm <sup>-3</sup>	O.M. (%)	Al	Ca	Mg	H + Al	CEC	V	Al Sat. (%)	K	S	B	Mn	Zn	Cu
											cmolc dm <sup>-3</sup>								
											mg dm <sup>-3</sup>								
LU1	40	4.7	4.9	12	102	6.4	1.4	3.1	1.1	15.4	19.6	22	24	1.3	32	0.5	76	9.1	7.3
LU2	38	4.8	4.9	6	65	2.8	2.7	1.8	0.5	15.4	17.7	14	52	0.9	22	0.4	83	9.6	6.8

Cy Clay content; SPM=buffer solution; .OM = soil organic matter; CEC= Cation exchange capacity; V =base saturation Ca+Mg+K/S+Al); S = Ca+Mg+K, in cmolc kg<sup>-1</sup>

Fonte: Os Autores.

**Figure 2.** Species accumulation curves using the Bootstrap estimator. LU1: Landscape Unit 1 (low anthropogenic pressure); LU2: Landscape Unit 2 (high anthropogenic pressure)..



Fonte: Os Autores.

### ***INFLUENCE OF HUMAN ACTION ON OCCURRENCE OF MACROPHYTE SPECIES***

The plant richness in LU2 was restricted to occurrence of four species, with highest occurrence frequency of *A. philoxeroides*, indicating that the phytosociological conditions were favorable for its development. This dominance of *A. philoxeroides* can be justified by conditions such as: high adaptation of the species to low soil fertility and exclusion of other species by selective grazing pressure, since the biomass of the species is little affected by fragmentation (Dong et al. 2012). *Alternanthera philoxeroides* was found from the flooded to the driest environment, with wide distribution and high degree of soil cover, with value 8 in 47% and 9 in 53% of the quadrats assessed on the van der Maarel scale. In LU2, the colonization and competition potential of this species, native to South America, was confirmed, for being herbaceous perennial, stoloniferous, with pronounced clonal propagation (Kissmann & Groth 2000, Gunasekera & Bonila 2001). The clonal habit is a common propagation strategy among the macrophytes, which together with the high phenotypic plasticity, make this species invasive in other environments (Ward et al. 2008, Geng et al. 2007).

The adaptation of *A. philoxeroides* in LU2 influenced the occurrence of other species negatively, e.g., those found in the LU1. The species *Paspalum notatum* cf was recorded exclusively in LU2 with the second highest occurrence frequency (Table 1), but had a low soil cover degree in the quadrats. The genus *Paspalum* is well-adapted to wetlands, has forage value and almost all species are perennial (Penteado & Macedo 2000, Rocha & Lins 2009, Maciel et al. 2010). The fact of being associated with nutrient-poor soils can explain its occurrence in LU2 (Table 3). In LU2, the rudimentary agriculture and extensive grazing may have contributed to the selection of this species by

selective grazing of animals, resulting in reduced species richness in LU2 (Table1). It was not possible to evaluate the total number of species in LU1 (Figure 2). In this landscape occurred greater wealth of species and also of life forms related to different habitats (floating leaves, emergent vegetative part and amphibious – Table 1). The evaluation in LU2 shows tendency of stabilization of the accumulation curve, indicating that probably all the species were evaluated. Carvalho et al. 2011 support the hypothesis of a positive relationship between community resilience and richness. Even with potential species to collect, this study supports this hypothesis.

In tropical regions, the low human impact stimulates a greater diversity of macrophytes which is desirable in flooded environments, which are responsible for the receipt, retention and export of sediments and nutrients. Such areas also mitigate climate changes, represent a biodiversity reserve and can be used for tourism, recreation and appreciation of the natural and cultural heritage.

## CONCLUSION

The damming over 50 years ago allowed an investigation of two areas surrounding the reservoir, where it was found that that most procumbent amphibious native macrophytes species are located in areas that are less affected by humans. A greater protection of the areas surrounding the reservoir, in other words, a reduced human impact, allows a greater diversity of macrophytes. This response is desirable in this environment type, for functioning as a buffer zone and avoiding pollutants entry. The intensification of land use in the reservoir surroundings restricted macrophyte diversity.

## ACKNOWLEDGEMENTS

The authors wish to thank the undergraduate students of biology for their valuable help with the field work. E.C. is indebted to the Brazilian Council for Scientific and Technological Development (CNPq) for a research productivity grant (CNPq, Pq 2).

## REFERENCES

- Aitkenhead-Peterson JA, McDowell WH, Neff JC 2003. Sources, Production and Regulation of Allochthonous Dissolved Organic Matter Inputs to Surface. In: Findlay SEG, Sinsabaugh RL, eds. *Aquatic ecosystems Interactivity of dissolved organic matter*. San Diego, Elsevier/Academic Press, p.26-59.
- Alves JA, Tavares AS, Trevisan R 2011. Composição e distribuição de macrófitas aquáticas na lagoa da Restinga do Massiambú. Área de Proteção Ambiental Entorno Costeiro, SC. *Rodriguésia* 62(4):785-801.
- Bailey-Serres J, Voesenek LACJ 2008. Flooding stress: acclimations and genetic diversity. *Annu Rev Plant Biol* 59:313-339.
- Banach K, Banach AM, Lamers LPM, Kroon H, Bennicelli RP, Smits AJM, Visser EJW 2009. Differences in flooding tolerance between species from two wetland habitats with contrasting

- hydrology: implications for vegetation development in future floodwater retention areas. *Ann Bot* 103:341–351.
- Boldrini II, Trevisan R, Schneider AA 2008. Estudo florístico e fitossociológico de uma área às margens da lagoa do Armazém, Osório, Rio Grande do Sul, Brasil. *Rev Bras Bioc* 6(4):355-367.
- Brasil 2005. *Resolução Nº 357, de 17 de março de 2005 do Conselho Nacional de Meio Ambiente*.
- Braun-Blanquet J, Capdevilla J, Oriol B. 1979. *Fitosociologia: bases para el estudio de las comunidades vegetales*. Madrid, H. Blume, 820 pp.
- Carvalho P, Thomaz SG, Kobayashi JT, Bini LM 2012. Species richness increases the resilience of wetland plant communities in a tropical floodplain. *Austral Ecol* 38(5):592-598.
- CEEE – Companhia Estadual de Geração e Transmissão de Energia Elétrica 2011. *Plano de Uso e Ocupação do Entorno dos Reservatórios da UHE Capigui*. Porto Alegre, Profill Engenharia e Ambiente 54 pp.
- Clesceri LS, Greenberg AE, Eaton AD 1999. *Standart Methods for examination of water and freshwater, 20th edn*. Washington, American Public Health Association 541 pp.
- Colmer TD, Voeselek LACJ. 2009. Flooding tolerance: suites of plant traits in variable environments. *Funct Plant Biol* 36:665- 681.
- Colwell RK 2013. EstimateS: Statistical estimation of species richness and shared species from samples. User's Guide and application. Avialbe from: <http://purl.oclc.org/estimates>
- Colwell RK, Coddington JA 1994. Estimating terrestrial biodiversity through extrapolation. *Philos Trans R Soc Lond B Biol Sci* 345(1311):101–118.
- CQFS-RS/SC 2016. *Manual de adubação e calagem para os estados do Rio Grande do Sul e de Santa Catarina*, 10 ed. Porto Alegre: Sociedade Brasileira de Ciência do Solo, Comissão de Química e Fertilidade do Solo Rio Grande do Sul e de Santa Catarina, 376 pp.
- Cunha GR 1997. *Meteorologia: fatos & mitos*. Passo Fundo, Embrapa/CNPT, 267 pp.
- Chambers PA, Lacoul P, Murphy KJ, Thomaz SM 2008. Global diversity of aquatic macrophytes in freshwater. *Hydrobiology* 595:9-26.
- Dong B-C, Alpert P, Guo W, Yu F-H 2012. Effects of fragmentation on the survival and growth of the invasive, clonal plant *Alternanthera philoxeroides*. *Biol Invasions* 14:1101–1110.
- Evangelista HBA, Thomaz SM, Evangelista LR 2012. Comparison of Diversity Indices Applied to Macrophyte. *Braz Arch Biol Technol* 55(2):277-282.
- Evans DE 2003. Aerenchyma formation. *New phytol* 161:35-49.
- Ferreira FA, Mormul RP, Thomaz SM, Pott A, Pott VJ 2011. Macrophytes in the upper Paraná river floodplain: checklist and comparison with other large South American wetlands. *Rev Biol Trop* 59(2):541-556.
- Galvão AKL, Silva JF, Albertino SMF, Monteiro GFP, Cavalcante DP 2011. Levantamento fitossociológico em pastagens de várzea no estado do Amazonas. *Planta Daninha* 29(1):69-75.

- Geng YP, Pan XY, Xu C, Zhang WJ, Li B, Chen JK 2007. Phenotypic plasticity rather than locally adapted ecotypes allows the invasive alligator weed to colonize a wide range of habitats. *Biol Invasions* 9:245–256.
- Irgang BE, Gastal JCVS 1996. *Macrófitas aquáticas da planície costeira do Rio Grande do Sul*. Porto Alegre, CPG-Botânica/UFRGS, 290 pp.
- IUSS Working Group WRB 2014. *World reference base for soil resources 2014. International soil classification system for naming soils and creating legends for soil maps*. World Soil Resources Reports FAO, Rome, Italy 106 pp.
- Jackson MB, Colmer TD 2005. Response and adaptation by plants flooding stress. *Ann Bot* 96:501-505.
- Jackson MB, Ishizawa K, Ito O 2009. Evolution and mechanisms of plant tolerance to flooding stress. *Ann Bot* 103:37-142.
- Kelman J. Pereira, MVF, Neto TAA, Sales PRH, Viera AM 2006. Hidreletricidade. In: Rebouças AC, Braga B, Tundisi JG (Org.) *Águas doces no Brasil, capital ecológico, uso e conservação*, 3 ed. São Paulo, Escrituras, p. 507-543.
- List of flora species of Brazil. Jardim Botânico do Rio de Janeiro. [updated 29 dez 2016] Available from <http://floradobrasil.jbrj.gov.br/>
- Maarel EVD 1979. Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* 39(2):97-114.
- Maciel JR, Oliveira RC, Alves M 2010. Aspectos da distribuição e conservação de *Paspalum L.* (Poaceae) em Pernambuco. *Hoehnea* 37(4):801-812.
- Maëmets H, Palmik K, Sudnitsuna D, Melnik M 2010. Eutrophication and macrophyte species richness in the large shallow North-European Lake Peipsi. *Aquat Bot* 92:273–280.
- Maltchik L. Rolon AS, Stenert C, Machado IF, Rocha O 2011. Can rice field channels contribute to biodiversity conservation in Southern Brazilian wetlands? *Rev Biol Trop* 59(4):1895-1914.
- Meyer ST, Franceschinelli EV 2010. Estudo florístico de plantas vasculares associadas às áreas úmidas na cadeia do Espinheiro (MG), Brasil. *Rev Bras Bot* 33(4):677-691.
- Mukhopadhyay G, Dewanji A 2005. Presence of tropical hydrophytes in relation to limnological parameters - a study of two freshwater ponds in Kolkata, India. *Ann Limnol* 41(4):281-289.
- Navarro-Ortega A, Acuña V, Bellin A, Burek P, Cassiani G, Redouane CA, Dolédec A, Elosegi A, Ferrari F, Ginebreda A, Grathwohl P, Jones C, Rault PK, Kok K, Koundouri P, Ludwig RP, Merz R, Milacic R, Barceló D 2015. Managing the effects of multiple stressors on aquatic ecosystems under water scarcity. The GLOBAQUA project. *Sci Tot Environ* 503–504:3-9.
- OECD 2015. Governança dos Recursos Hídricos no Brasil, [ebook on the internet]. OECD Publishing, Paris. Available from: <http://dx.doi.org/10.1787/9789264238169-pt>
- Paron LM, Freitas DH, Pereira CM 2011. *Manual de Procedimentos de amostragem e análise físico química da água (recurso eletrônico)*. Embrapa Florestal, Colombo, 67 pp.

- Peintinger M, Pratib D, Winkler E 2007. Water level fluctuations and dynamics of amphibious plants at Lake Constance: Long-term study and simulation Perspect. *Plant Ecol* 81:79-196.
- Penteado MIO, Macedo MCM 2000. Avaliação de espécies forrageiras em áreas mal drenadas dos cerrados. *Rev Bras Zoot* 29(6):1616-1622.
- Pereira AS, Cláudio RIT, Edélti FA, Cleber PS 2012. Aquatic macrophytes as indicators of water quality in subtropical shallow lakes Southern Brazil. *Acta Limnol Bras* 24:52-53.
- Pereira SA 2012. Aquatic macrophytes as indicators of water quality in subtropical shallow lakes Southern Brazil. *Acta Limnol Bras* 24:52-53.
- Pott VJ, Pott A, Lima LCP, Moreira SN, Oliveira AKM 2011. Aquatic macrophyte diversity of the Pantanal wetland and upper basin. *Braz Jour of Biol* 71(1):255-263.
- Rocha AES, Lins ALA 2009. Checklist das Poaceae de áreas inundáveis e inundadas do nordeste do estado do Pará. *Acta Amazon* 39(4):763-772.
- Rolon AS, Rocha O, Maltchik L 2011. Diversity of aquatic macrophytes in the Lagoa do Peixe National Park. *Neotrop Biol Conserv* 6(1):5-12.
- Sairam RK, Kumutha D, Ezhilmathi K, Deshmukh PS, Srivastava GC 2008. Physiology and biochemistry of waterlogging tolerance in plants. *Biol Plant* 52(3):401-412.
- Sass LL, Bozek MA, Hauxwell JA, Wagner K, Knigh S 2010. Response of aquatic macrophytes to human land use perturbations in the watersheds of Wisconsin lakes, U.S.A. *Aquat Bot* 93:1-8.
- Spellmeier J, Périco E, Freitas EM 2009. Composição florística de um banhado no município de Estrela/Rio Grande do Sul. *Pesquisas: Série Botânica* 60:367-381.
- Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ 1995. Análise de solo, plantas e outros materiais, 2 ed. Porto Alegre, Editora da Universidade Federal do Rio Grande do Sul. 174 pp.
- Ward SM, Gaskin JF, Wilson LM 2008. Ecological Genetics of Plant Invasion: What Do We Know? *Invasive Plant Sci Manag* 1:98-109.
- Wetzel RG 2001. *Limnology Lake and river ecosystems* 3rd edn. San Diego, Elsevier Academic Press, 387 pp.
- Zhao Q, Liu S, Deng L, Dong S, Wang C. 2014. Soil degradation with water-level fluctuations in the Manwan Reservoir, Lancang River Basin. *Catena* 113:33-43.

## Ocorrência de Macrófitas como Resposta da Pressão Antropogênica em Reservatório

### RESUMO

O objetivo deste estudo foi o de avaliar o efeito do uso antrópico do solo sobre a comunidade de plantas aquáticas no entorno de um reservatório artificial consolidado. Foram delimitadas duas unidades paisagísticas (UP) no entorno de um reservatório artificial de mais de 50 anos: a UP1 apresenta baixo impacto antrópico e a UP2 apresenta alto impacto antrópico. No total, foram encontradas 20 espécies de plantas. Apenas duas espécies foram comuns em UP1 e UP2. A UP1 apresentou maior diversidade de espécies (n=18). Os teores de fósforo, potássio, matéria orgânica, cálcio e magnésio foram menores na UP2 (n=4). Pode-se concluir que o processo de barramento permite a ocorrência de maior número de espécies e de plantas de macrófitas, pela conservação do solo adjacente ao reservatório. O baixo impacto antrópico nessa área permite maior diversidade de macrófitas nativas. A intensificação do uso do solo no entorno do reservatório não permite o desenvolvimento de comunidade diversificada de macrófitas, pois empobrece o solo.

**Palavras-chave:** Plantas Aquáticas; Conservação; Áreas Úmidas.

Submission: 29/12/2016

Acceptance: 04/12/2017